

Vitamin D Concentrations and Obstructive Sleep Apnea in a Multicenter Cohort of Older Males

Umesh Goswami¹, Kristine E. Ensrud^{1,2}, Misti L. Paudel², Susan Redline^{3,4}, Eva S. Schernhammer^{3,4}, James M. Shikany⁵, Katie L. Stone⁶, Ken M. Kunisaki^{2,1} for the Osteoporotic Fractures in Men (MrOS) Study Research Group.

1. University of Minnesota, Minneapolis, MN, US
2. Minneapolis VA Health Care System, Minneapolis, MN, US
3. Brigham and Women's Hospital, Boston, MA, US
4. Harvard Medical School, Boston, MA, US
5. University of Alabama at Birmingham, Birmingham, AL, US
6. California Pacific Medical Center, San Francisco, CA, US

Corresponding Author:

Ken Kunisaki, MD, MS
Minneapolis VA Health Care System
Pulmonary, Critical Care and Sleep (111N)
1 Veterans Drive
Minneapolis, MN, USA, 55417
kunis001@umn.edu

Word Count: 3189

Descriptor Number: 8.28

Running Title: Vitamin D and obstructive sleep apnea

Keywords (MeSH terms): Sleep apnea, obstructive; Vitamin D; Obesity; Cross-Sectional Study.

Author Contributions: Umesh Goswami: contributed to the study design, data collection, data analysis, manuscript writing, and final editing. He had full access to all the data in the study and takes responsibility for the integrity and accuracy of the data analysis. Kristine E. Ensrud: contributed to the study design, data analysis, manuscript writing, and final editing. Misti L. Paudel: contributed to the study design, data collection, data analysis, manuscript writing, and final editing. Susan Redline: contributed to the study design, manuscript writing, and final editing. Eva S. Schernhammer: contributed to the study design, manuscript writing, and final editing. James M. Shikany: contributed to the study design, manuscript writing, and final editing. Katie L. Stone: contributed to the study design, manuscript writing, and final editing. Ken M. Kunisaki: contributed to the study design, data collection, data analysis, manuscript writing, and final editing. He had full access to all the data in the study and takes responsibility for the integrity and accuracy of the data analysis.

Funding Support: The Osteoporotic Fractures in Men (MrOS) Study is supported by National Institutes of Health funding. The following institutes provide support: the National Institute of

Arthritis and Musculoskeletal and Skin Diseases (NIAMS), the National Institute on Aging (NIA), the National Center for Research Resources (NCRR), and NIH Roadmap for Medical Research under the following grant numbers: U01 AR45580, U01 AR45614, U01 AR45632, U01 AR45647, U01 AR45654, U01 AR45583, U01 AG18197, U01 AG027810, and UL1 TR000128. The National Heart, Lung, and Blood Institute (NHLBI) provides funding for the MrOS Sleep ancillary study "Outcomes of Sleep Disorders in Older Men" under the following grant numbers: R01 HL071194, R01 HL070848, R01 HL070847, R01 HL070842, R01 HL070841, R01 HL070837, R01 HL070838, and R01 HL070839. Funding for the Vitamin D assays was provided under the grant number: R01 AG030089. The Veterans Health Administration Office of Research and Development also provided protected research time in support of this study.

Previous Presentation: Some of these data were previously presented as a poster abstract at the American Thoracic Society International Conference, San Diego, May 2014.

Disclaimer: The views expressed in this article are those of the authors and do not reflect the views of the U.S. Government, the Department of Veterans Affairs, or any of the authors' affiliated institutions.

Abstract

Rationale: Seasonal nadirs in 25-hydroxyvitamin D (25[OH]D) concentrations overlap with increased incidence and severity of obstructive sleep apnea (OSA) in winter. We hypothesized that because lower 25(OH)D concentrations might lead to upper airway muscle dysfunction, low 25(OH)D would be associated with higher apnea-hypopnea index (AHI), a measure of OSA severity.

Objectives: To determine if lower 25(OH)D concentration is associated with greater prevalence and increased severity of OSA, independent of established OSA risk factors.

Methods: Using unconditional logistic regression, we performed a cross-sectional analysis in the Outcomes of Sleep Disorders in Older Men study which included in-home overnight polysomnography, serum 25(OH)D measurement, and collection of demographic and co-morbidity data. The primary outcome was severe sleep apnea as defined by AHI \geq 30/hr.

Measurements and Main Results: Among 2,827 community-dwelling, largely Caucasian (92.2%), elderly ([mean \pm SD] age 76.4 \pm 5.5 years) males, mean 25(OH)D concentration was 28.8 \pm 8.8 ng/mL. Subjects within the lowest quartile of 25(OH) D (6-23 ng/mL) had greater odds of severe sleep apnea in unadjusted analyses (OR 1.45; 95% CI: 1.02-2.07) when compared to the highest 25(OH)D quartile (35-84 ng/mL). However, further adjustment for established OSA risk factors strongly attenuated this association (multivariable adjusted OR 1.05; 95% CI: 0.72-1.52), with body mass index and neck circumference as the main confounders. There was also no evidence of an independent association between lower 25(OH)D levels and increased odds of mild (AHI 5.0-14.9/hr) or moderate (AHI 15.0-29.9/hr) sleep apnea.

Conclusions: Among community-dwelling older men, the association between lower 25(OH)D and sleep apnea was largely explained by confounding by larger BMI and neck circumference.

Abstract Word Count: 257

Obstructive sleep apnea (OSA) is a disease of recurrent, partial or complete upper airway closure during sleep. The pathogenesis of OSA is complex¹ and a recent study reported increased OSA incidence and severity during winter months². Although the magnitude of the effect was modest (median difference in apnea-hypopnea index [AHI] of 3.3 events/hr), the authors hypothesized that winter-related fat redistribution, medication use, fluid displacement to the neck, and/or air pollution were the reasons for their observation. However, an alternative explanation of the seasonal variation in OSA may be the coinciding winter nadir of vitamin D³.

Low circulating 25-hydroxyvitamin D [25(OH)D] concentrations are associated with poor musculoskeletal function^{4,5}. Control of upper airway muscle tone is felt to be a major contributor to OSA¹ and therefore, patients with low 25(OH)D concentrations might have an increased risk of OSA due to worse function of the skeletal muscle supporting upper airway patency during sleep. Low 25(OH)D concentrations are also associated with airway inflammation, chronic rhinitis and repeated upper airway infections leading to tonsillar enlargement⁶⁻¹⁰, which may additionally contribute to OSA incidence and severity. Low 25(OH)D concentrations are also associated with type 2 diabetes mellitus, metabolic syndrome and obesity, all of which are frequently found in patients with OSA^{11,12}. Only a few studies have studied the association between lower 25(OH)D levels and OSA and these have reported inconsistent results¹³⁻¹⁵.

To better address this knowledge gap, we analyzed data from a large, multi-center, community-based cohort study and tested the hypothesis that lower blood concentrations of 25(OH)D are associated with greater prevalence and increased severity of OSA, independent of

classic OSA risk factors.

Methods

Study Participants

The Osteoporotic Fractures in Men (MrOS) Study enrolled 5,995 community-dwelling men aged 65 and older during the baseline examination between 2000 and 2002^{16,17}. To be eligible for the study, men had to be able to walk without assistance and not have had a bilateral hip replacement. Participants were recruited at six clinical centers (Birmingham, AL; Minneapolis, MN; Palo Alto, CA; Monongahela Valley near Pittsburgh, PA; Portland, OR; and San Diego, CA). The Outcomes of Sleep Disorders in Older Men Study (MrOS Sleep) visit occurred on average 3.4 ± 0.5 years (range 1.9–4.9) after the baseline examination, between December 2003 and March 2005. Ethics approval was obtained from the institutional review board at each site and the Coordinating Center and Reading Center. Written informed consent for participation in the MrOS Sleep Study was obtained for all individuals.

The MrOS Sleep Study was an ancillary study with a target recruitment number of 3,000 men from the parent MrOS Study. Exclusions for the MrOS Sleep Study included use of nocturnal positive airway pressure or oral appliance devices, use of supplemental oxygen use, and presence of an open tracheostomy. Of the 5,995 MrOS participants, 3,135 participated in the MrOS Sleep Study, while 2,860 participants in the main cohort did not participate in the MrOS Sleep Study. Of the non-participants in the MrOS Sleep Study, 1,997 refused participation, and compared to those who participated in MrOS Sleep, those who refused

participation were older by 1 year (74.01 years \pm 5.88 vs. 73.05 years \pm 5.55, $p < 0.001$) and not significantly different with respect to body mass index (BMI; 27.20 \pm 3.72 kg/m² vs. 27.38 \pm 3.72 kg/m², $p = 0.10$)¹⁸.

Of 3,135 MrOS Sleep Study participants, 2,911 (92.8 %) had at least 4 hours of technically adequate sleep study data for analysis, and of these, 2,827 (90.2 %) had 25(OH)D concentrations measured. These 2,827 men constituted the analytical cohort for the current study.

Polysomnography and other sleep-related measures

In-home sleep studies using unattended polysomnography (Safiro, Compumedics, Inc., Melbourne, Australia) were performed. The recording montage consisted of C3/A2 and C4/A1 electroencephalograms, bilateral electrooculograms, a bipolar submental electromyogram, thoracic and abdominal respiratory inductance plethysmography, airflow (using nasal-oral thermocouple and nasal pressure cannula), finger pulse oximetry, electrocardiogram, body position (mercury switch sensor), and bilateral leg movements (piezoelectric sensors).

Trained certified staff members performed home visits for setup of the sleep study units. After sensors were placed and calibrated, signal quality and impedance were checked, and sensors were repositioned as needed to improve signal quality, replacing electrodes if impedances were greater than 5 k Ω , using approaches similar to those in the Sleep Heart Health Study¹⁹. After studies were downloaded, they were transferred to the Case Reading Center (Cleveland, OH) for centralized scoring by a trained technician. Polysomnography data quality was excellent, with a failure rate of less than 4% and more than 70% of studies graded

as being of excellent or outstanding quality. Quality codes for signals and studies were graded using previously described approaches, which included coding the duration of artifact-free data per channel and overall study quality (reflecting the combination of grades for each channel). We note that central apnea events were included in the overall apnea-hypopnea index (AHI), but such events were rare (only 7% of study participants had a central apnea index ≥ 5 /hr).

Vitamin D analysis

Serum for vitamin D analysis was collected at baseline of the MrOS Sleep Study and immediately frozen at -70°C . Concentrations of 25-hydroxyvitamin D2 and 25-hydroxyvitamin D3 were measured by liquid chromatography-tandem mass spectrometry (LC-MS/MS) (ThermoFisher Scientific, Franklin, Massachusetts 02038 and Applied Biosystems-MDS Sciex, Foster City, CA 94404) at the Mayo Clinic Reference Laboratories (Singh RJ, PhD, Mayo Clinic Laboratory, Rochester, MN), using fasting samples collected at the Sleep Visit. Deuterated stable isotope (d3-25-hydroxyvitamin D) was added to a 0.2 ml serum sample as internal standard. 25-hydroxyvitamin D2, 25-hydroxyvitamin D3, and the internal standard were extracted using acetonitrile precipitation.

The extracts were then further purified online and analyzed by LC-MS/MS using multiple reaction monitoring. Using three different target markers as quality controls for each assay, inter-assay CVs for 25-hydroxyvitamin D3 were 9.7% at 9.0 IUs, 7.5% at 29 IUs, and 5.8% at 76 IUs. For 25-hydroxyvitamin D2, CVs were 11.2% at 11 IUs, 8.5% at 28 IUs, and 7.7% at 74 IUs. The minimum detectable limit for 25-hydroxyvitamin D2 was 4 ng/ml and for 25-hydroxyvitamin D3 was 2 ng/ml. Total 25-hydroxyvitamin D [25(OH)D] was calculated by

adding 25-hydroxyvitamin D2 and 25-hydroxyvitamin D3.

Clinical data

All covariate data were collected at the time of the sleep study visit. All participants completed questionnaire data, which included questions about medical history, smoking, and alcohol intake. Hypertension was defined as a positive response to the question, “Has a doctor or other healthcare provider told you that you have hypertension or high blood pressure?” Race was based on self-report and categorized as Caucasian, African American, Asian, or Hispanic/other. BMI was calculated as weight (kg)/height (m²), and obesity was defined as a BMI greater than 30 kg/m². During the home or clinic visits, body weight was measured using a standard balance beam scale and height using a wall-mounted Harpenden stadiometer (Holtain, UK). Neck and waist circumference were also measured using standard methods. Snoring was assessed according to self-report.

The MacArthur Subjective Status Scale (range 1–10) was used to assess perceived social status, with higher scores representing higher perceived social status. Participants were asked to bring in all current medications used within the preceding 30 days. All prescription and non-prescription medications were entered into an electronic database and each medication was matched to its ingredient(s) based on the Iowa Drug Information Service (IDIS) Drug Vocabulary (College of Pharmacy, University of Iowa, Iowa City, IA). A variable for season (January-March=Winter; April-June=Spring; July-September=Summer; October-December=Fall) was calculated using the date of the participant’s clinic exam.

Statistical Analysis

In primary analyses, we expressed 25(OH)D concentrations in quartiles and compared baseline characteristics across quartiles using analysis of variance or chi-square testing for continuous or categorical variables, respectively. We also used the following fixed 25(OH)D categories in our analyses: <20 ng/mL, 20-29.9 ng/mL, and ≥ 30 ng/mL; but because results were similar, we present quartiles as our primary 25(OH)D variable.

We used unconditional logistic regression models to calculate odds ratios and 95% confidence intervals for 25(OH)D quartiles (referent group quartile 4) for the primary dichotomous outcome of severe sleep apnea as defined by $\text{AHI} \geq 30/\text{hr}$ and a secondary dichotomous outcome of at least moderate sleep apnea defined by $\text{AHI} \geq 15/\text{hr}$. We created four logistic regression models including an unadjusted model, a model adjusted for established OSA risk factors^{18, 20-22} (age, BMI, neck circumference and hypertension), a model further adjusted for season of blood draw²³ and a fully adjusted model that included the above variables along with medications that may affect upper airway patency (opiates, benzodiazepines and alcohol)^{20, 24, 25}, race, smoking and clinic site. We conducted a sensitivity analysis to evaluate the effect of selected variables (BMI, neck circumference, hypertension) on the outcome by adding one variable at a time to the crude model. We also conducted two secondary analyses expressing 25(OH)D using clinical cutpoints (<20 ng/mL, 20-29.9 ng/mL, and ≥ 30 ng/mL)²⁶ and a polytomous regression model comparing outcomes of $\text{AHI} 0\text{-}4.9/\text{hr}$ (no OSA) to standard OSA severity categories of mild ($\text{AHI} 5\text{-}14.9/\text{hr}$), moderate ($\text{AHI} 15\text{-}29.9/\text{hr}$) and severe ($\text{AHI} \geq 30/\text{hr}$) OSA.

In exploratory analyses, to examine effect modification by BMI, we evaluated the

association between 25(OH)D and OSA excluding men with BMI ≥ 30 kg/m² hypothesizing that the relationship between 25(OH)D and OSA would be stronger in non-obese men. We also tested for an interaction between 25(OH)D and BMI for the prediction of severe OSA, using the log-likelihood ratio test.

Results

Among 3,135 participants who attended the MrOS Sleep visit, 2,911 (92.9 %) participants had at least 4 hours of technically adequate sleep study data for analysis. 2,827 (90.2 %) of these men had 25(OH)D concentrations measured and constituted the analytical cohort.

The study cohort consisted of largely Caucasian (92.2%) older men ([mean \pm SD] age 76.4 \pm 5.5 years), with a mean BMI of 27.2 \pm 3.8 kg/m². The demographic and co-morbid disease characteristics of the study participants are provided in Table 1.

The mean 25(OH)D concentration was 28.8 \pm 8.8 ng/mL, with concentrations ≥ 30 ng/mL (widely considered to represent replete vitamin D status) in 1,247 (44.1%), between 20-29.9 ng/mL (widely considered vitamin D insufficient) in 1,205 (42.6%), and < 20 ng/mL (widely considered vitamin D deficient) in 375 (13.3%). The distribution of AHI categories 0-4.9, 5-14.9, 15-29.9 and ≥ 30 events/hour was 1,105 (39%), 977 (34.6%), 470 (16.6%) and 276 (9.8%) respectively. Among the 276 participants with AHI ≥ 30 /hr, 50 (18.1%) were 25(OH)D deficient (< 20 ng/mL), compared to 134 (12.1%) who were 25(OH)D deficient among those with AHI < 5 /hr ($p=0.06$).

Participants in the lowest quartile of 25(OH) D (6-23 ng/mL) had greater odds of AHI ≥ 30 /hr (crude OR 1.45; 95% CI 1.02-2.07), compared to participants in the highest 25(OH)D quartile (35-84 ng/mL) (Table 2). After adjustment for traditional OSA risk factors, this association was no longer evident (adjusted OR 1.05; 95%CI 0.72-1.52). Further adjustment for season and other covariates did not alter these results. Findings were similar when AHI ≥ 15 /hour was substituted for AHI ≥ 30 /hr, when 25(OH)D was expressed using clinical cutpoints of <20 ng/mL, 20-29.9 ng/mL, and ≥ 30 ng/mL, and in polytomous regression evaluating the association between 25(OH)D quartiles and odds of no OSA vs. mild/moderate/severe OSA (Table 3).

Sensitivity analysis suggested that the association between lower 25(OH)D concentrations and higher odds of OSA was largely explained by greater BMI and larger neck circumference among those men with lower 25(OH)D concentrations (Table 4).

In exploratory analysis restricted to those with BMI <30 kg/m² (n=2,255; 79.8% of overall cohort), the modest association between low 25(OH)D and higher odds of severe sleep apnea did not reach significance (fully-adjusted OR [quartile 1 vs. quartile 4] of AHI ≥ 30 /hr = 1.27; 95%CI 0.82-1.97; fully-adjusted OR [25(OH)D <20 ng/mL vs. ≥ 30 ng/mL] of AHI ≥ 30 /hr = 1.44 ; 95% CI 0.90-2.28). There was no evidence of an interaction between obesity (BMI <30 vs. ≥ 30 kg/m²) and 25(OH)D for the prediction of AHI ≥ 30 /hr, when 25(OH)D was categorized by quartiles (p=0.197) or by clinical cutpoints (p=0.562).

Discussion

Despite plausible mechanisms potentially connecting vitamin D deficiency to OSA pathogenesis and severity, we found no evidence of an independent association between 25(OH)D concentration and OSA in our analyses of this cohort of older community-dwelling men.

Our findings suggest that the association between lower 25(OH)D levels and a higher odds of OSA was due in large part to greater BMI and larger neck circumference among participants with lower 25(OH)D levels. These results indicate that low 25(OH)D may simply be a marker of larger BMI and neck circumference, rather than directly contributing to OSA pathogenesis.

Several lines of evidence link obesity with lower 25(OH)D concentrations. Obese subjects are more likely to be restricted in physical activity, thus limiting their exposure to sunlight and resulting in lower 25(OH)D concentrations²⁷. Additionally, inflammatory cytokines upregulated in adiposity are known to inversely affect 25(OH)D bioavailability and increase its metabolic clearance²⁸. Finally, poor dietary habits leading to obesity often provide a poor source of oral vitamin D intake.

We explored the possibility that 25(OH)D effects on OSA pathogenesis might be most pronounced in non-obese participants, since mechanisms other than soft tissue accumulation and external airway pressure might account for loss of upper airway patency during sleep in non-obese individuals. In these analyses, we observed similar non-significant findings, although the adjusted OR point estimates for low 25(OH)D and AHI \geq 30/hr were higher than in analyses that included both obese and non-obese individuals. There was no evidence of an interaction between obesity and 25(OH)D on OSA, but further studies with larger sample sizes are needed

to more definitively evaluate the potential role of vitamin D deficiency in the pathogenesis of OSA in non-obese individuals.

Despite reasons to believe that vitamin D deficiency may play a role in the pathogenesis of OSA, few studies to date have explored this potential association and these have reported mixed results. In a selected population of 150 adult OSA patients (50 patients each with mild, moderate and severe OSA) and controls matched on BMI, gender and age (n=32), Mete and colleagues¹³ noted no significant difference in 25(OH)D concentrations between OSA patients and controls ($17.9 \pm 9.3 \mu\text{g/dL}$ vs. $19.2 \pm 7.2 \mu\text{g/dL}$; $p=0.468$). However, the authors reported that among those with $\text{AHI} >30/\text{hr}$, 78% were 25(OH)D deficient ($<20 \mu\text{g/dL}$), compared to 50% among controls with $\text{AHI} <5/\text{hr}$ ($p=0.02$). We found similar results in our population of older men, although our observed difference in prevalence of 25(OH)D deficiency was smaller in magnitude (18.1% among those with $\text{AHI} \geq 30/\text{hr}$, and 12.1% among those with $\text{AHI} <5/\text{hr}$).

However, the study by Mete and colleagues notably preselected obese patients in their study (mean BMI of 32 kg/m^2), and they were therefore not able to adjust for the effect of BMI on the relationship between 25(OH)D and AHI. We have addressed this in our study by enrolling an unselected population of community-dwelling older men with a wide range of BMI (and AHI) and adjusting our analysis for BMI and several other potential confounders.

In contrast to the results of Mete and colleagues and our study, Kheirandish-Gozal and colleagues¹⁴ found a statistically significant correlation between 25(OH)D concentrations and AHI ($r = -0.285$, $p < 0.001$) in pediatric OSA patients (mean age 6.5 to 7.2 years). The relevance of these results to adult OSA is not clear, because pediatric OSA has a very different pathogenesis than adult OSA. In children, upper airway inflammation, in the form of

adenotonsillar hypertrophy, is felt to be the major contributor to OSA²⁹ as opposed to adults, in whom fat redistribution, upper airway muscle dysfunction and age-related changes in upper airway anatomy are felt to be the major contributors¹.

More recently, Bertisch and colleagues¹⁵ studied the association between 25(OH)D concentrations and various sleep measures including AHI in a multi-center, multi-ethnic cohort of 1,721 adults with a mean age of 68.2 years. The authors reported that those with 25(OH)D <20 ng/mL had a statistically higher median AHI (2.1 events/hr higher) than those with 25(OH)D >29 ng/mL. Similar to our findings, those with lower 25(OH)D concentrations were also more obese. Also similar to our findings, the AHI difference was no longer significant after adjusting for covariates such as age, gender, and waist circumference.

In exploratory analyses, the authors found that AHI was higher by 7.1 events/hour in Chinese-American participants, (n=205) with low 25(OH)D compared to higher 25(OH)D, but this was not found in the Caucasian, African-American or Hispanic-American participants. The authors importantly noted that 25(OH)D concentrations were measured an average of 10.3 years prior to the collection of polysomnography data, so whether or not 10-year old 25(OH)D measures reflect 25(OH)D status at the time of polysomnography is a major limitation. Our 25(OH)D and polysomnography data were concurrent and therefore truly cross-sectional. Nevertheless, the data from their cohort and our cohort would appear to be consistent with an overall lack of association between 25(OH)D concentrations and sleep apnea.

The main strengths of our study are the large sample size and the community-based sample of older men. Our participants were not preselected for presence of any condition, particularly OSA, vitamin D deficiency, or obesity, thus minimizing selection biases. Despite the

community-based sampling design, we had a wide distribution of AHI, 25(OH)D concentrations, and BMI, thus allowing us to analyze such relationships across a wide range of these variables seen in clinical practice. Additional strengths of our study include performance of 25(OH)D assays at an experienced, high-quality reference laboratory, careful review and cleaning of sleep study data, and accurate measurement of potential confounders of 25(OH)D and OSA.

Our study has some inherent limitations as well. Our study participants were generally healthy, largely Caucasian, elderly males, so we cannot generalize these findings to non-Caucasians, females and younger patients. This was also a cross-sectional analysis with a single night's measurement of sleep and a single 25(OH)D measurement. Although our data suggest that low 25(OH)D is not likely to predict future development of OSA, a longitudinal study design would be required to specifically test such a hypothesis. We also note that although measures of OSA typically display high night-to-night reliability³⁰, 25(OH)D levels vary by season, latitude, skin tone, sunscreen use, and time spent outdoors. We adjusted for season in our regression analysis, but we had no measures of within-participant seasonal variation in 25(OH)D concentrations or other factors that may have varied over time. Therefore, our single 25(OH)D measure may not fully reflect seasonal variations throughout the year.

The aim of our study was also to assess vitamin D status irrespective of vitamin D source (outdoor sunlight exposure, diet, or oral supplement use). Our analyses were adjusted for season of blood draw and geographic region, but we acknowledge that different sources of vitamin D may have differential impacts on factors such as seasonal variations in 25(OH)D concentrations and prevalence of individuals with vitamin D deficiency. Lastly, we note that some data suggest that myopathy related to vitamin D deficiency may be most pronounced at

very low 25(OH)D concentrations such as <12 ng/mL³¹. Because our community-dwelling cohort had very few persons with such low levels (only 2.1% had concentrations <12 ng/mL), our study is not able to adequately address whether profound vitamin D deficiency could be independently associated with OSA.

Conclusions

Among community-dwelling older men, the association between lower serum 25(OH)D concentrations and higher odds of sleep apnea was explained by greater BMI and larger neck circumference among those with lower 25(OH)D levels.

Acknowledgements

We thank the participants in the MrOS Sleep study.

Table 1. Characteristics of 2827 participants by 25(OH)D quartiles

Variable	Overall Cohort (n=2827)	Serum 25(OH) D Quartiles				P Value
		Q1 6-23 ng/mL (n=767)	Q2 24-28 ng/mL (n=686)	Q3 29-34 ng/mL (n=678)	Q4 35-84 ng/mL (n=696)	
Age, years	76.4 ± 5.5	76.8 ± 5.6	76.4 ± 5.5	76.1 ± 5.4	76.3 ± 5.6	0.10
Caucasian, %	92.2	88.7	92.4	93.7	94.4	0.0002
BMI, kg/m ²	27.2 ± 3.8	28.1 ± 4.2	27.5 ± 4.0	26.7 ± 3.4	26.3 ± 3.1	<.0001
Neck circumference, cm	39.4 ± 2.8	40.0 ± 2.9	39.6 ± 2.9	39.2 ± 2.6	38.9 ± 2.6	<.0001
Systolic blood pressure, mmHg	126.8±16.2	126.8±16.9	126.7±16.1	126.8±16.3	126.9±15.7	0.99
Current Smoker, %	2.0	2.4	2.2	1.2	2.2	0.68
Alcohol intake, > 14 drinks/wk, %	5.4	3.9	6.0	3.6	8.2	<.0001
Benzodiazepine use, %	4.63	4.82	4.52	4.28	4.89	0.95
Opiate use, %	3.30	5.22	3.64	3.10	3.30	0.137
Winter (Jan-March), %	33.4	42.6	32.1	31.0	26.7	<.0001
Spring (April-June), %	25.3	26.6	27.1	26.3	21.0	<.0001
Summer (July-Sept), %	22.6	15.8	24.1	23.0	28.2	<.0001
Fall (October-Dec), %	18.8	15.0	16.8	19.8	24.1	<.0001
AHI 0-4.9/hr, %	39.1	37.2	39.7	39.1	40.5	0.26
AHI 5-14.9/hr, %	34.6	34.9	31.3	36.3	35.6	0.26
AHI 15-29.9/hr, %	16.6	16.8	17.8	15.9	16.0	0.26
AHI ≥30/hr, %	9.8	11.1	11.2	8.7	7.9	0.26
AHI, events/hr	11.8 ± 12.9	12.4 ± 13.3	12.4 ± 13.2	11.4 ± 12.7	10.9 ± 12.5	0.08
25(OH)D, ng/mL	28.8 ± 8.8	18.5 ± 4.0	26.0 ± 1.4	31.4 ± 1.7	40.37 ± 5.5	<.0001

Table 2: Logistic regression for odds of obstructive sleep apnea

25(OH)D Quartiles	OR (95% CI) of AHI ≥15/hr	OR (95% CI) of AHI ≥30/hr
Crude		
Q1 vs. Q4	1.24 (0.98-1.56)	1.45 (1.02-2.07)
Q2 vs. Q4	1.31 (1.03-1.66)	1.47 (1.02-2.12)
Q3 vs. Q4	1.04 (0.82-1.34)	1.11 (0.76-1.63)
p-for-trend	0.027	0.016
Adjusted for typical OSA risk factors*		
Q1 vs. Q4	0.96 (0.75-1.23)	1.05 (0.72-1.52)
Q2 vs. Q4	1.11 (0.87-1.43)	1.18 (0.81-1.72)
Q3 vs. Q4	0.98 (0.76-1.27)	1.06 (0.72-1.57)
p-for-trend	0.940	0.737
Adjusted for typical OSA risk factors* and 25(OH)D modifiers [#]		
Q1 vs. Q4	0.94 (0.72-1.22)	0.97 (0.65-1.44)
Q2 vs. Q4	1.13 (0.87-1.46)	1.15 (0.78-1.70)
Q3 vs. Q4	1.00 (0.77-1.29)	1.05 (0.71-1.56)
p-for-trend	0.827	0.931
Fully adjusted [^]		
Q1 vs. Q4	0.94 (0.72-1.23)	0.96 (0.65-1.43)
Q2 vs. Q4	1.13 (0.87-1.47)	1.15 (0.78-1.70)
Q3 vs. Q4	0.98 (0.75-1.27)	0.99 (0.66-1.48)
p-for-trend	0.884	0.993

*: "typical OSA risk factors" covariates: age, body mass index, neck circumference, hypertension

#: winter season: January – March

[^]: fully adjusted covariates: age, body mass index, neck circumference, hypertension, winter season, clinic site, race, alcohol consumption, smoking, benzodiazepine use, opioid use

25(OH)D = 25-hydroxyvitamin D

AHI = apnea-hypopnea index

CI = confidence interval

OR = odds ratio

OSA = obstructive sleep apnea

Table 3: Polytomous logistic regression for odds of obstructive sleep apnea

25(OH)D Quartiles	OR (95% CI)		
	AHI 5-14.9/hr vs. AHI 0-4.9/hr	AHI 15-29.9/hr vs. AHI 0-4.9/hr	AHI ≥30/hr vs. AHI 0- 4.9/hr
Crude			
Q1 vs. Q4	1.07 (0.84-1.36)	1.15 (0.85-1.56)	1.53 (1.05-2.23)
Q2 vs. Q4	0.90 (0.70-1.15)	1.14 (0.84-1.55)	1.45 (0.99-2.13)
Q3 vs. Q4	1.06 (0.83-1.35)	1.04 (0.76-1.42)	1.14 (0.76-1.71)
p-for-trend	0.073		
Adjusted for typical OSA risk factors*			
Q1 vs. Q4	0.89 (0.69-1.14)	0.87 (0.64-1.19)	0.97 (0.65-1.45)
Q2 vs. Q4	0.80 (0.62-1.03)	0.96 (0.70-1.32)	1.07 (0.72-1.60)
Q3 vs. Q4	1.00 (0.78-1.28)	0.95 (0.69-1.31)	1.05 (0.69-1.59)
p-for-trend	0.534		
Adjusted for typical OSA risk factors* and 25(OH)D modifiers#			
Q1 vs. Q4	0.91 (0.70-1.18)	0.88 (0.63-1.24)	0.91 (0.59-1.39)
Q2 vs. Q4	0.82 (0.63-1.06)	0.99 (0.72-1.38)	1.06 (0.70-1.61)
Q3 vs. Q4	1.01 (0.79-1.30)	0.98 (0.70-1.36)	1.05 (0.69-1.60)
p-for-trend	0.720		
Fully adjusted^			
Q1 vs. Q4	0.91 (0.70-1.19)	0.89 (0.63-1.25)	0.90 (0.59-1.38)
Q2 vs. Q4	0.81 (0.62-1.06)	0.99 (0.71-1.38)	1.06 (0.70-1.61)
Q3 vs. Q4	0.99 (0.7-1.28)	0.98 (0.70-1.36)	0.98 (0.64-1.50)
p-for-trend	0.739		

*: "typical OSA risk factors" covariates: age, body mass index, neck circumference, hypertension

#: winter season: January – March

^: fully adjusted covariates: age, body mass index, neck circumference, hypertension, winter season, clinic site, race, alcohol consumption, smoking, benzodiazepine use, opioid use

25(OH)D = 25-hydroxyvitamin D

AHI = apnea-hypopnea index

CI = confidence interval

OR = odds ratio

OSA = obstructive sleep apnea

Table 4. Sensitivity Analysis. Logistic regression with adjustment for only one typical OSA risk factor (age, BMI, neck circumference, hypertension).

25(OH)D Quartiles	OR (95% confidence Intervals)	
	AHI \geq 15 vs. <15	AHI \geq 30 vs. <30
Adjusted for continuous age		
Q1 vs. Q4	1.22 (0.97-1.55)	1.43 (1.00-2.04)
Q2 vs. Q4	1.30 (1.02-1.65)	1.47 (1.02-2.11)
Q3 vs. Q4	1.05 (0.82-1.34)	1.12 (0.77-1.65)
p-trend	0.035	0.024
Adjusted for continuous BMI		
Q1 vs. Q4	0.99 (0.78-1.27)	1.12 (0.77-1.61)
Q2 vs. Q4	1.14 (0.89-1.45)	1.23 (0.85-1.78)
Q3 vs. Q4	0.99 (0.77-1.27)	1.04 (0.70-1.53)
p-trend	0.806	0.429
Adjusted for neck circumference		
Q1 vs. Q4	1.07 (0.84-1.36)	1.22 (0.85-1.75)
Q2 vs. Q4	1.18 (0.92-1.50)	1.29 (0.89-1.87)
Q3 vs. Q4	0.99 (0.77-1.28)	1.07 (0.73-1.57)
p-trend	0.354	0.201
Adjusted for hypertension		
Q1 vs. Q4	1.23 (0.97-1.56)	1.45 (1.01-2.06)
Q2 vs. Q4	1.32 (1.04-1.68)	1.49 (1.03-2.14)
Q3 vs. Q4	1.04 (0.81-1.33)	1.11 (0.75-1.63)
p-trend	0.028	0.016

References

1. Dempsey JA, Veasey SC, Morgan BJ, O'Donnell CP. Pathophysiology of sleep apnea. *Physiol Rev.* 2010;90(1):47-112.
2. Cassol CM, Martinez D, da Silva FA, Fischer MK, Lenz Mdo C, Bos AJ. Is sleep apnea a winter disease? Meteorologic and sleep laboratory evidence collected over one decade. *Chest.* 2012;142(6):1499-1507.
3. van Schoor N, Knol D, Deeg D, Peters F, Heijboer A, Lips P. Longitudinal changes and seasonal variations in serum 25-hydroxyvitamin D levels in different age groups: results of the Longitudinal Aging Study Amsterdam. *Osteoporosis Int.* 2014;25(5):1483-91.
4. Bischoff-Ferrari HA, Dietrich T, Orav EJ, Hu FB, Zhang Y, Karlson EW, Dawson-Hughes B. Higher 25-hydroxyvitamin D concentrations are associated with better lower-extremity function in both active and inactive persons aged > or =60 y. *Am J Clin Nutr.* 2004 Sep;80(3):752-8
5. Janssen HC, Samson MM, Verhaar HJ. Vitamin D deficiency, muscle function, and falls in elderly people. *Am J Clin Nutr.* 2002;75(4):611-615.
6. Wang L, Lee C, Chien C, Chen JY, Chiang F, Tai C. Serum 25-hydroxyvitamin D levels are lower in chronic rhinosinusitis with nasal polyposis and are correlated with disease severity in Taiwanese patients. *Am J Rhinology Allergy.* 2013;27(6):e162-e165.
7. Abuzeid WM, Akbar NA, Zacharek MA. Vitamin D and chronic rhinitis. *Curr Opin Allergy Clin Immunol.* 2012;12(1):13-17.
8. Jung J, Kim J, Cho S, Choi B, Min K, Kang H. Allergic rhinitis and serum 25-hydroxyvitamin D level in Korean adults. *Ann Allergy Asthma Immunol.* 2013;111(5):352-357.
9. McNicholas W, Tarlo S, Cole P et al. Obstructive apneas during sleep in patients with seasonal allergic rhinitis. *Am Rev Respir Dis.* 1982;126(4):625.
10. Reid D, Morton R, Salkeld L, Bartley J. Vitamin D and tonsil disease--preliminary observations. *Int J Pediatr Otorhinolaryngol.* 2011;75(2):261-264.
11. Johnson T, Avery G, Byham-Gray L. Vitamin D and metabolic syndrome. *Top Clin Nutr.* 2009;24(1):47-54.
12. Bozkurt NC, Cakal E, Sahin M, Ozkaya EC, Firat H, Delibasi T. The relation of serum 25-hydroxyvitamin-D levels with severity of obstructive sleep apnea and glucose metabolism abnormalities. *Endocrine.* 2012;41(3):518-525.
13. Mete T, Yalcin Y, Berker D et al. Obstructive sleep apnea syndrome and its association with vitamin D deficiency. *J Endocrinol Invest.* 2013;36(9):681-5.
14. Kheirandish-Gozal L, Peris E, Gozal D. Vitamin D levels and obstructive sleep apnoea in children. *Sleep Med.* 2014;15(4):459-463.

15. Bertisch SM, Sillau S, de Boer IH, Szklo M, Redline S. 25-hydroxyvitamin D concentration and sleep duration and continuity: Multi-Ethnic Study of Atherosclerosis. *Sleep*. 2015;38(8):1305-1311.
16. Orwoll E, Blank JB, Barrett-Connor E et al. Design and baseline characteristics of the osteoporotic fractures in men (MrOS) study—a large observational study of the determinants of fracture in older men. *Contemp Clin Trials*. 2005;26(5):569-585.
17. Blank JB, Cawthon PM, Carrion-Petersen ML et al. Overview of recruitment for the osteoporotic fractures in men study (MrOS). *Contemp Clin Trials*. 2005;26(5):557-568.
18. Mehra R, Stone KL, Blackwell T et al. Prevalence and Correlates of Sleep-Disordered Breathing in Older Men: Osteoporotic Fractures in Men Sleep Study. *J Am Geriatr Soc*. 2007;55(9):1356-1364.
19. Redline S, Sanders MH, Lind BK et al. Methods for obtaining and analyzing unattended polysomnography data for a multicenter study. Sleep Heart Health Research Group. *Sleep*. 1998;21(7):759-767.
20. Stradling JR, Crosby JH. Predictors and prevalence of obstructive sleep apnoea and snoring in 1001 middle aged men. *Thorax*. 1991;46(2):85-90.
21. Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med*. 1993;328(17):1230-1235.
22. Young T, Skatrud J, Peppard PE. Risk factors for obstructive sleep apnea in adults. *JAMA*. 2004;291(16):2013-2016.
23. Kasahara AK, Singh RJ, Noymer A. Vitamin D (25OHD) Serum Seasonality in the United States. *PloS one*. 2013;8(6):e65785.
24. Webster LR, Choi Y, Desai H, Webster L, Grant BJ. Sleep-Disordered Breathing and Chronic Opioid Therapy. *Pain Med*. 2008;9(4):425-432.
25. Berry RB, Kouchi K, Bower J, Prosser G, Light RW. Triazolam in patients with obstructive sleep apnea. *Am J Respir Crit Care Med*. 1995;151(2 Pt 1):450-454.
26. LeFevre ML, U.S. Preventive Services Task Force. Screening for vitamin D deficiency in adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2015;162(2):133-140.
27. Parikh SJ, Edelman M, Uwaifo GI et al. The relationship between obesity and serum 1, 25-dihydroxy vitamin D concentrations in healthy adults. *J Clin Endocrinol Metab*. 2004;89(3):1196-1199.
28. Wortsman J, Matsuoka LY, Chen TC, Lu Z, Holick MF. Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr*. 2000;72(3):690-693.
29. Marcus CL. Sleep-disordered breathing in children. *Am J Respir Crit Care Med*. 2001;164(1):16-30.

30. Redline S, Tosteson T, Boucher M, Millman R. Measurement of sleep-related breathing disturbances in epidemiologic studies. Assessment of the validity and reproducibility of a portable monitoring device. *Chest*. 1991;100(5):1281-1286.
31. Bischoff HA, Stahelin HB, Urscheler N et al. Muscle strength in the elderly: its relation to vitamin D metabolites. *Arch Phys Med Rehabil*. 1999;80(1):54-58.